

Healthy Blue — Medicaid UM Guideline

Subject: Autism Spectrum Disorder Services
Status: Active

Current Effective Date: 07/01/2017
Last Review Date: 8/9/2023

Description

Adaptive behavioral treatment (ABT) described as intensive behavioral intervention (IBI) or applied behavioral analysis (ABA) for autism spectrum disorder (ASD). In this document, the term 'ABT,' which includes services such as ABA and IBI, refers to services provided as part of ABA and IBI.

Services to treat ASD, as defined in the most recent edition of the Diagnostic and Statistical Manual of Mental Disorders, are provided to eligible Medicaid beneficiaries 0 to 21. ASD services must be recommended by a physician, Licensed Psychologist or a Licensed Psycho-Educational Specialist (LPES) within his or her scope of practice under South Carolina state law to prevent the progression of ASD, prolong life, and promote the physical and mental health and efficiency of the individual. These services may be provided in the beneficiary's home, clinical setting or other settings as authorized.

A licensed psychologist, developmental pediatrician or a Licensed Psycho-Educational Specialist (LPES) certified by the South Carolina Department of Education to perform such evaluations, and acting within the scope of their competency, must certify and document through a comprehensive psychological assessment or testing report that the beneficiary meets the medical necessity criteria for services via a DSM ASD diagnosis. The comprehensive psychological testing assessment or report should include:

1. A clinical interview with the beneficiary or family members or guardians as appropriate.
- 2. A review of the presenting problems, symptoms and functional deficits, strengths and history, including past psychological assessment reports and records.
- 3. Assessments also include a behavioral observation in one or more settings.
- 4. ASD diagnosis from current edition of DSM, including severity levels
- 5. Autism Diagnostic Observation Schedule (ADOS; ADOS-2).
- 6. At least one of the following:
 - Autism Diagnosis Interview (ADI)
 - Behavior Assessment System for Children (BASC)
 - Childhood Autism Rating Scale (CARS)
 - Gilliam Autism Rating Scale (GARS)
 - Vineland Adaptive Behavioral Scales (Vineland)
 - Social Responsiveness Scale (SRS)
 - A standardized measure of intelligence (e.g., WISC or WAIS, Stanford-Binet, Bayley Scales, etc.)
 - Screening checklists (e.g., MCHAT, STAT, ASQ, etc.)

** This link leads to a third-party site. That organization is solely responsible for the contents and privacy policies on its site.

www.HealthyBlueSC.com

BlueChoice HealthPlan is an independent licensee of the Blue Cross Blue Shield Association.

To report fraud, call our confidential Fraud Hotline at **800-763-0703**. You may also call the South Carolina Department of Health and Human Services Fraud Hotline at **888-364-3224** or email fraudres@scdhhs.gov.

BlueChoice HealthPlan-Medicaid UM Guideline

Subject: Autism Spectrum Disorder Services

Current Effective Date: 07/01/2017

Status: Active

Last Review Date: 06/11/2021

- Social Communication Questionnaire (SCQ)

ASD Treatment Services by a LIP:

ASD treatment services can only be furnished by a licensed independent practitioner (LIP) via prior authorization once an Individualized Plan Of Care (IPOC) has been submitted and approved. ASD treatment services are evidence-based practices (EBP) that support the amelioration and management of symptoms specific to the diagnosis of ASD. Direct beneficiary contact (and collaterals as clinically indicated) are necessary for billable ASD Treatment Services.

Allowable EBPs for ASD treatment services by a LIP include:

- Cognitive Behavioral Intervention Package (CBIP)
- Comprehensive Behavioral Treatment for Young Children (CBTYC)
- Language Training (Production)
- Modeling
- Naturalistic Teaching Strategies (NTS)
- Parent Training Package
- Peer Training Package
- Pivotal Response Treatment®
- Schedules
- Scripting
- Self-Management
- Social Skills Package
- Story-based Interventions

ASD Treatment Services by an Applied Behavior Analysis Provider:

Assessments are to be completed by a Board Certified Behavior Analyst (BCBA) or Board Certified Assistance Behavior Analyst (BCaBA). Assessments may include direct observation and measurement of beneficiary behavior in structured and unstructured situations, determination of baseline levels of adaptive and maladaptive behaviors and functional behavior analysis.

Assessment for IPOC Development

Behavior Identification Assessment:

A behavior identification assessment is an initial functional assessment to be completed by a BCBA or a BCaBA under the supervision of a BCBA. The behavior identification assessment includes direct beneficiary contact (and collaterals as clinically indicated) in order to identify maladaptive behaviors, complete a mental health evaluation and establish treatment needs and an IPOC.

This service must include at least two of the following assessments:

- Vineland Adaptive Behavior Scales (Vineland)
- Verbal Behavior Milestone Assessment and Placement Program (VBMAPP)

BlueChoice HealthPlan-Medicaid UM Guideline

Subject: Autism Spectrum Disorder Services

Current Effective Date: 07/01/2017

Status: Active

Last Review Date: 06/11/2021

- Peabody Picture Vocabulary Test
- Social Responsiveness Scale
- Assessment of Functional Living Skills (AFLS)
- Essentials for Living (EFL)
- Assessment of Basic Language and Learning

Skills (ABBLs)

Treatment services for ABA are provided via a prior authorization once an IPOC is submitted and approved. ABA services are furnished by a BCBA-D, a BCBA, a BCaBA or an RBT in accordance with their competency parameters, as per the Behavior Analyst Certification Board (BCAB). BCaBAs and RBTs furnishing services must be under the direction supervision of a BCBA within their scope of competency.

ABA treatment services include:

1. Observational behavioral follow-up assessment — direct beneficiary contact (and collaterals as clinically indicated) to identify and evaluate factors that may impede adaptive behavior. This service includes a functional assessment of problem behavior based on direct observation data psychological testing, as clinically indicated.
2. Adaptive behavior treatment by protocol — direct beneficiary contact (and collaterals as clinically indicated) to address the beneficiary's treatment goals
3. Adaptive behavior treatment with protocol modification — direct beneficiary contact to resolve one or more factors impeding adaptive behavior and may simultaneously instruct a technician and or family or caregiver in the modified treatment goals
4. Family adaptive behavior treatment guidance — direct contact with the family or caregiver for specialized training and education to assist with the beneficiary's treatment goals and development. The provider observes and trains the family or caregivers on the beneficiary's status, as well as instructs family or caregivers on techniques to promote the child's development.
5. Group adaptive behavior treatment by protocol — direct contact with two or more beneficiaries to address the beneficiary's treatment goals

Clinical Indications

Medically Necessary:

Treatment plan signed and submitted by licensed MD/DO/NP/PA which must include the following required elements:

- Clinical interview with the member or family or caregiver.
- Completed psychological testing results performed by a licensed psychologist, developmental pediatrician or a LPES certified by the South Carolina Department of Education to perform such evaluations, and acting within the scope of their competency.
- DSM-5-TR criteria met for autism including severity scale.
- Autism Diagnostic Observation Schedule (ADOS; ADOS-2).

BlueChoice HealthPlan-Medicaid UM Guideline

Subject: Autism Spectrum Disorder Services

Current Effective Date: 07/01/2017

Status: Active

Last Review Date: 06/11/2021

- One of the following standard psychiatric assessments:
 - Behavior Assessment System for Children (BASC).
 - Vineland Adaptive Scale (within 95 percent confidence interval).
 - Autism Diagnostic Interview-Revised (ADI-R).
 - Social Communications Questionnaire (SCQ).
 - Childhood Autism Rating Scale (CARS).
 - Autism Behavior Checklist (ABC).
 - Modified Checklist for Autism in Toddlers (M-CHAT).
 - Gilliam Autism Rating Scale (GARS).
- BCBA/BCaBA assessment and program plan (IPOC) that must include two of the following elements:
 - ABLLS, AFLS, EFL, VB-MAPP, Vineland, Peabody Picture Vocabulary Test, SCQ.
 - Behavioral observation in one or more settings, presenting problems, symptoms and functional deficits, anticipated outcomes stated as measurable goals related to each specific problematic behavior or skill deficit (language, imitation, social skills, cooperation, etc.).
 - Achievement timeframes.
 - Submission of signed FCM-287 ABA Service Provider Request form.

Recertification Requests

Treatment plan signed and submitted by licensed MD which must include the following required elements:

- BCBA/BCaBA assessment and program plan that must include the following elements:
 - ABLLS, AFLS, EFL or VB-MAPP.
 - Anticipated outcomes stated as measurable goals related to each specific problematic behavior or skill deficit (language, imitation, social skills, cooperation, etc.). These will be submitted on a new form currently in development.
 - Achievement timeframes.
 - Amount and type of parent or caregiver participation, as applicable to member.
 - Date of every ninety-day review and annual re-development.
 - Signature, title and date by the multidisciplinary team members including the parent or caregiver.
 - Submission of signed FCM287-ABA Service Provider Request form.
 - Depending on the benefit structure, providers may be asked to submit the last two 90-Day Progress Summaries including the following information: specific objective(s) towards which the 90 days has focused, specific treatment activities and interventions, goals that have been met, graphs of goals and objective demonstrating progress, explanation of delayed progress toward goals, amount and time of parent/caregiver participation, summary of treatment plan for the upcoming 90-days and signature and title, and date by the multidisciplinary team members to include the parent and/or caregiver (see Attachment A for examples of graphs).

The IPOC must be completed no later than:

- The tenth business day after an initial assessment meeting with a LIP or a behavior assessment with an ABA provider is completed; **OR**
- If the IPOC is not completed within this time frame, services rendered are not Medicaid reimbursable.

Note: The number of hours allotted for direct treatment with the individual can continue to be up to 40 hours a week. The hours should be reviewed regularly, and adjusted to address the behavioral

BlueChoice HealthPlan-Medicaid UM Guideline

Subject: Autism Spectrum Disorder Services
Status: Active

Current Effective Date: 07/01/2017
Last Review Date: 06/11/2021

targets and key functional skills of the individual, based on the results of the assessments previously mentioned.

* Systematic and repeated evaluation of developmental status is critical to assessing the effect of therapeutic treatments, including ABT. The use of standardized assessments facilitates the consistent, systematic, and reliable evaluation early in the course of treatment, preferably before initiating ABT, and at regularly scheduled intervals thereafter. The data derived from these assessments is used to inform about the impact of treatment on the trajectory of the individual's condition, especially documenting improvement.

Reporting/Documentation

The clinical record must contain documentation sufficient to justify Medicaid reimbursement and should allow an individual not familiar with the beneficiary to evaluate the course of treatment.

The beneficiary's clinical record must include, at a minimum, the following documentation:

- A comprehensive assessment or testing report which establishes medical necessity via an ASD diagnosis.
- A Behavioral Identification Assessment for ABA services.
- Signed, titled and dated IPOC.
- Signed releases, consents, Beneficiary Rights acknowledgment and confidentiality assurances for treatment.
- Signed, titled and dated clinical service note (CSN) and progress summaries.
- Copies of all written reports, and any other documents relevant to the care and treatment of the beneficiary.
- Consent for treatment that is dated and signed by the beneficiary, parent, legal guardian or primary care giver or legal representative. A new consent should be signed and dated with each authorization.
- Transition or discharge planning.
- Coordination of care.
- Emergency safety interventions.

**All ASD treatment services must be documented in CSNs within five (5) business days of their delivery. ABA providers must document in accordance with ABA standards and guidelines. All other providers' CSNs must include the nature of the beneficiary's treatment, any changes in treatment, discharge, crisis interventions and any changes in medical, behavioral or psychiatric status. Documentation must justify the amount of reimbursement claimed to Medicaid.

At minimum, all CSN's requires documentation of the following elements:

- Date.
- Face to face time start and end time.
- Individuals present during the visit.
- Brief description of services provided.
- Clinical note on the recipient's behavior.
- Place of service and delivery setting.

BlueChoice HealthPlan-Medicaid UM Guideline

Subject: Autism Spectrum Disorder Services

Current Effective Date: 07/01/2017

Status: Active

Last Review Date: 06/11/2021

- Any communication with guardians or caregivers.
- Signature of rendering provider with title.
- All elements must be documented in legible handwriting.
- Corrections to the medical record must adhere to the following guidelines:
- For paper record corrections draw one line through the error, and write “error”, “ER”, “mistaken entry” or “ME” to the side of the error in parenthesis. Enter the correction, sign or initial, and date it.
- Errors cannot be totally marked through; the information in error must remain legible. No correction fluid may be used.
- For electronic health records, error correction must include date or time stamp and user ID.
- Late entries should rarely be used, and then only used to correct a genuine error of omission or to add new information that was not discovered until a later date. Whenever late entries are made adhere to the following:
 - Identify the new entry as a “late entry.”
 - Include date and time and identify or refer to the date and incident for which late entry is written.

The IPOC must be individualized and specify problems to be addressed, goals to be worked toward and the strengths of the beneficiary. The IPOC must be developed prior to the delivery of a service with the full participation of the beneficiary and his or her family. The IOP must be completed in its entirety and include the following:

- Beneficiary’s strengths, needs, abilities and preferences.
- Goals and objectives of treatment which tie into the assessment and evaluation results.
- Outline to address the assessment needs.
- Specific treatment activities and interventions.
- Amount and type of parent or caregiver participation, as applicable.
- Date of each completed progress summary and annual re-development.
- Signature, title and date by the multidisciplinary team members including parent or caregiver.

**The IPOC must be completed no later than the 10th business day after an initial assessment meeting. If the IPOC is not completed within this timeframe, services rendered are not Medicaid reimbursable. The IPOC must be reviewed as part of the regular progress summary and the progress summaries must be completed at least quarterly. A new IPOC must be developed every 12 months.

BlueChoice HealthPlan-Medicaid UM Guideline

Subject: Autism Spectrum Disorder Services

Current Effective Date: 07/01/2017

Status: Active

Last Review Date: 06/11/2021

Coding

Billable Codes²:

All Medicaid beneficiaries must be ages 0 to 21 and have an established ASD diagnosis to meet medical necessity criteria in order to receive ASD treatment services.

A licensed psychologist or a school psychologist certified by the South Carolina Department of Education to perform such evaluations and acting within the scope of their competency must certify and document through a comprehensive psychological assessment or testing report that the beneficiary meets the medical necessity criteria for services via a DSM or ICD-10 ASD diagnosis. A prior-established diagnosis is also acceptable provided the comprehensive assessment or testing report adheres to the medical necessity guidelines. The following guidelines shall be used to determine medical necessity:

² Procedure and HCPC Codes provided by the state of South Carolina

BlueChoice HealthPlan-Medicaid UM Guideline

Subject: Autism Spectrum Disorder Services

Current Effective Date: 07/01/2017

Status: Active

Last Review Date: 06/11/2021

Service Name	Procedure Code	Qualifications	Frequency Limits
Screening and Diagnostic Assessment Services			
Behavior Identification Assessment	97151	BCBA-D BCBA BCaBA	32 units annually
ASD Treatment Services			
Adaptive Behavior Treatment by Protocol	97153	BCBA-D BCBA BCaBA RBT	160 units per week (in any combination)
Group Adaptive Behavior Treatment by Protocol	97154	BCBA-D BCBA BCaBA RBT	2-6 patients, up to 6 hrs/day
Adaptive Behavior Treatment with Protocol Modifications	97155	BCBA-D BCBA BCaBA	To be rendered at the rate of 10% of weekly therapy hours, up to 64 units per month (in any combination)
Family Adaptive Behavior Treatment Guidanc	97156	BCBA-D BCBA BCaBA	96 units annually (up to 24 hours a year)
Group Adaptive Behavior Treatment by Protocol	97158	BCBA-D BCBA BCaBA	2 – 6 patients, up to 6 hours a day
Non-ABA Treatment Services	H2019	Licensed Psychologist Licensed Psychoeducational Specialist Licensed Independent Social Worker, Clinical Practice Licensed Marriage and Family Therapist Licensed Professional Counselor	4 units per week

BlueChoice HealthPlan-Medicaid UM Guideline

Subject: Autism Spectrum Disorder Services
Status: Active

Current Effective Date: 07/01/2017
Last Review Date: 06/11/2021

Discussion/General Information

The clinical approach to a parent's initial complaints about developmental abnormalities is dictated by age. Parents may bring the child in to the physician as young as 12 months, though generally the first contact is around 24 months or later. With children under 24 months, it will be more problematic to arrive at correct and stable diagnosis, delaying an accurate diagnosis. This serves to underline the importance of the physician's skills, knowledge and experience. Parents who already have a child with ASD are often sensitized to early symptoms and these children are often brought in at even earlier ages. The estimated risk for ASD in younger siblings in identified patients is five percent.

Practice parameters for screening: these require immediate further evaluation:

- No babbling by 12 months
- No gesturing by 12 months (pointing, waving bye-bye, etc.)
- No single words by 16 months
- No two-word spontaneous phrases by 24 months
- ANY loss of ANY language or social skills at ANY age (Filipek, Accardo, Baranck, 1999)

The evaluation of individuals who test positive on a specific test or meet the practice parameters should commence a diagnostic evaluation with a complete history and physical examination, performed by a clinician experienced in the evaluation of ASD, in an attempt to determine a working diagnosis. This should include a complete history of the pregnancy, labor and delivery, exposure to toxins, drugs, infections and delivery complications. A full developmental history of all milestones of infant development also needs to be obtained, which would include the assessment of regression for previously learned abilities.

Further testing may include, but not be limited to:

- Vision and audiometric screening.
- IQ evaluation.
- Speech and language evaluation.
- Blood work: CBC, thyroid.
- Other labs (as indicated): metabolic studies (amino and organic acids, ammonia, lactate, etc.), cytogenetic studies (high resolution chromosome studies, Fragile X testing, translocation/deletion syndromes, etc.).
- Imaging: MRI of brain (not typically indicated).
- EEG (sleep deprived may be preferred), especially with a history of seizure spectrum behavior.
- Consults with psychiatrist, neurologist, pediatrician, psychologist, speech, occupational or physical therapist, etc., as indicated.

Medical conditions associated with autistic-like behaviors include, but are not limited to:

- Seizure disorders.
- Hearing problems.
- Fetal alcohol syndrome.
- Trisomy 21.
- Goldenhar syndrome.
- Hypothyroidism.
- Phenylketonuria.
- Cornelia de Lange syndrome.

BlueChoice HealthPlan-Medicaid UM Guideline

Subject: Autism Spectrum Disorder Services

Current Effective Date: 07/01/2017

Status: Active

Last Review Date: 06/11/2021

- Neurofibromatosis.
- Tourette's syndrome.
- Dandy-Walker Syndrome.
- Myoclonic epilepsy of infancy (Dravet Syndrome).
- Doose syndrome: myoclonic atstatic seizures.

There is also an emerging literature that suggests international adoption, and perhaps family movement to another language or culture, may severely impact infants, toddlers and preschoolers in the early phases of cognitive language development, such that autistic like behaviors may occur. This may be referred to as a disruption of cognitive language development, as opposed to communicative language development.

Applied behavior analysis (ABA) for ASD includes efforts to extinguish negative behaviors, and to replace these with positive behaviors by improving skills, based upon Skinnerian concepts of conditioned responses. Another focus would be the development of the ability to generalize concepts already learned to novel situations. Aside from a concentration on the core symptoms, there is a focus on attention and initiation as well. All involve a structured environment, predictable routines, functional as opposed to standardized treatment, a transition plan and significant family involvement. At the initial evaluation, target symptoms are identified, with designated interventions. There is also a need to provide an assessment mechanism at specified intervals.

ABA services for young children are often referred to as early intensive behavioral intervention (EIBI). EIBI targets cognition, language, social skills, etc. These services have typically been provided to children ages 8 and under for a duration of two to three years. During an average week, these services may be provided for up to 40 hours. Although the literature is clear that 10 hours per week results in a less robust therapeutic response, it remains unclear as to the optimal number of hours per week. After these preschool years, services may be provided by the child's school, as well as in the child's home. For children above the age of 10 years, services requested may focus on the development of social skills. Much of the ABA research has focused on children in the age range 2-7.

Other Therapies for ASD

Other therapies for ASD include, but are not limited to: auditory integration therapy, facilitated communication, developmental individual-difference relationship-based (DIR/floortime) model, relationship development intervention, holding therapy, movement therapies, music therapy, pet therapy, psychoanalysis, Son Rise program, scotopic sensitivity training, sensory integration therapy, neurotherapy (EEG biofeedback), gluten and casein free diets, mega-vitamin therapy, chelation of heavy metals, anti-fungal drugs for presumed fungal infection and secretin administration. There is insufficient evidence to support efficacy for a few of these treatments and no evidence for the majority. These programs are not considered to be behavioral therapy.

Aversive Techniques

Behavioral change techniques should use reinforcement whenever possible with consideration given to the least restrictive techniques possible. When punishment is chosen for a problem behavior, the techniques employed must follow the ethical guidelines dictated by the Behavior Analyst Certification Board. The parent or guardian must provide written consent for use of any techniques that may be considered aversive.

BlueChoice HealthPlan-Medicaid UM Guideline

Subject: Autism Spectrum Disorder Services
Status: Active

Current Effective Date: 07/01/2017
Last Review Date: 06/11/2021

Definitions

Autism spectrum disorder refers to a disorder as defined in the most recent edition of the *Diagnostic and Statistical Manual of Mental Disorders* of the American Psychiatric Association. This disorder is best characterized as neurobiological disorder of uncertain etiology, with symptom onset in the first years of life. Heritability is polygenic and estimated to be 90 percent. The primary phenomenon in ASD is severe impairment in development of social skills. Other deficits that may be observed include failure to initiate play or social interaction, inability to generalize learned behavior to new situations, failure to share attention with others, poor sleep, temper tantrums, hyperactivity, etc.

Autism is characterized by a triad of deficits in social skills, communication and language abilities and impairments in imaginative play. Additionally, there are often stereotypic behaviors, restricted interests and activities. Significant numbers of patients also experience intellectual disability and, as one would expect, there is a relationship between cognition and response to treatment.

Applied behavioral analysis for ASD refers to behavioral modification using ABA techniques, or the Lovaas Method, to target cognition, language and social skills. Techniques used within this paradigm include discreet trial training, incidental teaching, pivotal response training and verbal behavioral intervention. ABA service for younger children is often referred to as early intensive behavioral intervention (EIBI).

Board Certified Behavior Analyst® (BCBA®) refers to the person who is certified by the BACB (Behavioral Analyst Certification Board) and who is responsible for the administration of appropriate assessment tools, plan or notebook development, implementation and oversight, as well as workshop and training and oversight of all staff, reviewing data, updating plan, and troubleshooting. This person also conducts six-month assessments that include the Assessment of Basic Language and Learning Skills (ABLLS)– Assessment of Functional Living Skills (AFLS), Essential for Living (EFL) or Verbal Behavior Milestones Assessment and Placement (VB-MAPP), and a summary of treatment plan goals met and unmet related to specific behaviors, including, but not limited to, language, imitation, social skills, and cooperation, as well as periodic re-assessments of progress and problems noted.

Board Certified Assistant Behavior Analyst® (BCaBA®) refers to the person skilled in ABA techniques and is responsible for assessing progress via weekly monitoring utilizing highly structured formats and data review. BCaBAs may delegate services previously listed to a BCaBA who has been credentialed and is receiving required supervision. Rendering providers must possess a four-year college degree and may or may not be certified as BCBA. This person ensures that a RBT is well trained and acts as a liaison between the family and treatment team.

Registered Behavior Technician™ (RBT™) refers to the person trained in ABA techniques who is responsible for working directly with the child and for following a pre-set program plan for implementing skill acquisition and behavior reduction procedures and recording program data.

Educational interventions: learning interventions that assist children with obtaining knowledge and communication through speech, sign language, writing and other methods and social skills. **Note:** Many benefit contracts exclude coverage for services that are educational in nature.

BlueChoice HealthPlan-Medicaid UM Guideline

Subject: Autism Spectrum Disorder Services
Status: Active

Current Effective Date: 07/01/2017
Last Review Date: 06/11/2021

References

Government Agency, Medical Society, and Other Authoritative Publications:

1. Aman, M.G. (2005). Treatment planning for patients with autism spectrum disorders. Journal of Clinical Psychiatry, 66 (Suppl. 10), 38-45.
2. American Academy of Pediatrics Committee on Children With Disabilities. (2001). The pediatrician's role in the diagnosis and management of autistic spectrum disorder in children. Pediatrics, 107 (5), 1221-1226.
3. American Psychiatric Association. (2013). Diagnostic and Statistical Manual of Mental Disorders-5. Washington, DC: Author.
4. Bailey, D. B., Hebbeler, K., Spiker, D., Scarborough, A., Mallik, S. & Nelson, L. (2005). Thirty-Six-Month Outcomes for Families of Children Who Have Disabilities and Participated in Early Intervention. Pediatrics, 116 (6), 1346-1352.
5. Baranek, G. T., David, F. J., Poe, M. D., Stone, W. L. & Watson, L. R. (2006). Sensory Experiences Questionnaire: discriminating sensory features in young children with autism, developmental delays, and typical development. Journal of Child Psychology and Psychiatry, 47 (6), 591-601.
6. Ben-Itzhack, E. & Zachor, D. A. (2007). The effects of intellectual functioning and autism severity on outcome of early behavioral intervention for children with autism. Research in Developmental Disabilities, 28(3), 287-303.
7. Bodfish, J. (2004). Treating the core features of autism; are we there yet? Mental Retardation and Developmental Disabilities Research Reviews, 10 (4), 318-326.
8. Bryson, S. E., Zwaigenbaum, L., McDermott, C., Rombough, V. & Brian, J. (2008). The Autism Observation Scale for Infants: scale development and reliability data. Journal of Autism and Developmental Disorders, 38 (4), 731-738.
9. Capo, L. C. (2001). Autism, employment and the role of occupational therapy. Work, 16 (3), 201-207.
10. Cohen, H., Amerine-Dickens, M. & Smith, T. (2006). Early Intensive Behavioral Treatment: Replication of the UCLA Model in a Community Setting. Developmental and Behavioral Pediatrics, 27 (2), S145-155.
11. De Bildt, A., Sytema, S., Ketelaars, C., Kraijer, D., Mulder, E., Volkmar, F., and Minderaa, R. (2004). Interrelationship between Autism Diagnostic Observation Schedule-Generic (ADOS-G), Autism Diagnostic Interview-Revised (ADI-R), and the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV-TR) Classification in Children and Adolescents with Mental Retardation. Journal of Autism and Developmental Disorders, 34 (2), 129-137. Available online at: http://www.aspires-relationships.com/Interrelationship_Between_Autism_Diagnostic_Observation_Schedule_Generic.pdf. Last viewed on October 25, 2013.

BlueChoice HealthPlan-Medicaid UM Guideline

Subject: Autism Spectrum Disorder Services

Current Effective Date: 07/01/2017

Status: Active

Last Review Date: 06/11/2021

12. Eikeseth, S. (April 2008). Outcome of comprehensive psycho-educational interventions for young children with autism. *Research in Developmental Disabilities*; epublished April 1, 2008. Available online at: <http://www.asaonline.org/resources/research/psychoeducational>. Last viewed on October 25, 2013.
13. Eldevik, S., Eikeseth, S., Jahr, E. & Smith, T. (2006). Effects of Low-Intensity Behavioral Treatment for Children with Autism and Mental Retardation. *Journal of Autism and Developmental Disorders*, 36 (2), 211-224.
14. Eldevik, S., Hastings, R., Hughes, J., Jahr, E., Eikeseth, S. and Cross, S. (2009). Meta-Analysis of Early Intensive Behavioral Intervention for Children with Autism. *Journal of Clinical Child and Adolescent Psychology*, 38 (3), 439-450.
15. Eikeseth, S., Smith, T., Jahr, E. & Eldevik, S. (2002). Intensive Behavioral Treatment at School for 4- to 7-Year-Old Children with Autism: A 1-Year Comparison Controlled Study. *Behavior Modification*, 26 (1), 49-68.
16. Eikeseth, S., Smith, T., Jahr, E. & Eldevik, S. (2007). Outcome for Children With Autism Who Began Intensive Behavioral Treatment Between Ages 4 and 7: A Comparison Controlled Study. *Behavior Modification*, 31 (3), 264-278.
17. Filipek, P. A., Accardo, P. J., Baranek, G. T., Cook, E. H., Dawson, G., Gordon, B., Gravel, J. S., Johnson, C. P., Kallen, R. J., Levy, S. E., Minshew, N. J., Prizant, B. M., Rapin, I., Rogers, S. J., Stone, W. L., Teplin, S. W., Tuchman, R. F. & Volkmar, F. R. (1999). The screening and diagnosis of autism spectrum disorders. *Journal of Autism and Developmental Disorders*, 29 (6), 439-484.
18. Filipek, P. A., Accardo, P. J., Ashwal, S., Baranek, G. T., Cook, Jr., E. H., Dawson, G., Gordon, B., Gravel, J. S., Johnson, C. P., Kallen, R. J., Levy, S. E., Minshew, N. J., Ozonoff, S., Prizant, B. M., Rapin, I., Rogers, S. J., Stone, W. L., Teplin, S. W., Tuchman, R. F. & Volkmar, F. R. (2000). Practice parameter: Screening and diagnosis of autism: Report of the Quality Standards Subcommittee of the American Academy of Neurology and the Child Neurology Society. *Neurology*, 55 (4), 468-479.
19. Filipek, P. A., Steinberg-Epstein, R. & Book, T. M. (2006). Intervention for Autistic Spectrum Disorders. *NeuroRx™: The Journal of the American Society for Experimental NeuroTherapeutics*, 3 (2), 207-216.
20. Findling, R. L. (2005). Pharmacologic treatment of behavioral syndromes in autism and pervasive developmental disorders. *Journal of Clinical Psychiatry*, 66 (Suppl. 10), 26-31.
21. Fombonne, E. (2005). Epidemiology of autistic disorder and other pervasive developmental disorders. *Journal of Clinical Psychiatry*, 66 (Suppl. 10), 3-8.
22. Francis, K. (2005). Autism interventions: a critical update. *Developmental Medicine and Child Neurology*, 47 (7), 493-499.
23. Geschwind, D. H. (2009). Advances in Autism. *Annual Review of Medicine*, 60, 367-380.
24. Glogowska, M., Roulstone, S., Peters, T. J., & Enderby, P. (2006). Early speech- and language-impaired children: linguistic, literacy, and social outcomes. *Developmental Medicine & Child*

BlueChoice HealthPlan-Medicaid UM Guideline

Subject: Autism Spectrum Disorder Services

Current Effective Date: 07/01/2017

Status: Active

Last Review Date: 06/11/2021

Neurology, 48 (06), 489-494.

25. Gotham, K., Risi, S., Pickles, A. & Lord, C. (2007). The Autism Diagnostic Observation Schedule: revised algorithms for improved diagnostic validity. Journal of Autism and Developmental Disorders, 37 (4), 613-627.
26. Gurney, J. G., McPheters, M. L., & Davis, M. M. (2006). Parental report of health conditions and health care use among children with and without autism: National Survey of Children's Health. Archives of Pediatrics and Adolescent Medicine, 160 (8), 825-830.
27. Harris, S. (2003). Functional assessment. Journal of Autism and Developmental Disorders, 33 (2), 233.
28. Hastings, R. (2003). Behavioral adjustment of siblings of children with autism engaged in applied behavior analysis early intervention programs: The moderating role of social support. Journal of Autism and Developmental Disorders, 33 (2), 141-150.
29. Hoehn, T. P. & Baumeister, A. A. (1994). A Critique of the Application of Sensory Integration Therapy to Children with Learning Disabilities. Journal of Learning Disabilities, 27 (6), 338-350.
30. Howes, O. D., Rogdaki, M., Findon, J. L., Wichers, R. H., Charman, T., King, B. H., Loth, E., McAlonan, G. M., McCracken, J. T., Parr, J. R., Povey, C., Santosh, P., Wallace, S., Simonoff, E., & Murphy, D. G. (2018). Autism spectrum disorder: Consensus guidelines on assessment, treatment and research from the British Association for Psychopharmacology. Journal of Psychopharmacology, 33 (1), 3-29.
31. Howlin, P. (2005). The effectiveness of interventions for children with autism. Journal of Neural Transmission, Suppl. (69), 101-119.
32. Howlin, P., Magiati, I. & Charman, T. (2009). Systematic Review of Early Intensive Behavioral Interventions for Children with Autism. American Journal on Intellectual and Developmental Disabilities, 114 (1), 23-41.
33. Hyman, S. L. & Levy, S. E. (2005). Introduction: Novel Therapies in Developmental Disabilities – Hope, Reason, and Evidence. Mental Retardation and Developmental Disabilities, 11 (2), 107-109.
34. Hyman, S. L., Levy, S. E., & Myers, S. M. (2020). Identification, Evaluation, and Management of Children With Autism Spectrum Disorder. Pediatrics, 145 (1). Available online at: [Identification, Evaluation, and Management of Children With Autism Spectrum Disorder | Pediatrics | American Academy of Pediatrics \(aap.org\).](#)
35. Iarocci, G. & McDonald, J. (2006). Sensory Integration and the Perceptual Experience of Persons with Autism. Journal of Autism and Developmental Disorders, 36 (1), 77-90.
36. Itzchak, E. B., Lahat, E., Burgin, R. & Zachor, A. D. (2008). Cognitive, behavior and intervention outcome in young children with autism. Research in Developmental Disabilities, 29 (5), 447-458.
37. Johnson-Ecker, C. L. & Parham, L. D. (2000). The evaluation of sensory processing: A validity study using contrasting groups. American Journal of Occupational Therapy, 54 (5), 494-503.

BlueChoice HealthPlan-Medicaid UM Guideline

Subject: Autism Spectrum Disorder Services
Status: Active

Current Effective Date: 07/01/2017
Last Review Date: 06/11/2021

38. Kasari, C., Freeman, S. & Paparell, T. (2006). Joint attention and symbolic play in young children with autism: a randomized controlled intervention study. Journal of Child Psychology and Psychiatry, *47* (6), 611-620.
39. Keen, D., Sigafoos, J., & Woodyatt, G. (2001). Replacing prelinguistic behaviors with functional communication. Journal of Autism and Developmental Disorders, *31* (4), 385-398.
40. Kleinman, J. M., Ventola, P. E., Pandey, J., Verbalis, A. D., Barton, M., Hodgson, S., Green, J., Dumont-Mathieu, T., Robins, D. L. & Fein, D (2008). Diagnostic stability in very young children with autism spectrum disorders. Journal of Autism and Developmental Disorders, *38* (4), 606-615.
41. Koegel, L. K. (2000). Interventions to facilitate communication in autism. Journal of Autism and Developmental Disorders, *30* (5), 383-391.
42. Legoff, D. & Sherman, M. (2006). Long-term outcome of social skills intervention based on interactive LEGO© play. Autism, *10* (4), 317-329.
43. Linderman, T. M. & Stewart, K. B. (1999). Sensory integrative-based occupational therapy and functional outcomes in young children with pervasive developmental disorders: A single-subject study. American Journal of Occupational Therapy, *53* (2), 207-213.
44. Lord, C. (2000). Commentary; achievements and future directions for intervention research in communication and autism spectrum disorders. Journal of Autism and Developmental Disorders, *30* (5), 393-398.
45. Mailloux, Z. (1990). An Overview of the Sensory Integration and Praxis Tests. American Journal of Occupational Therapy, *44* (7), 589-594.
46. Magiati, I., Charman, T. & Howlin, P. (2007). A two-year prospective follow-up study of community-based early intensive behavioural intervention and specialist nursery provision for children with autism spectrum disorders. Journal of Child Psychology and Psychiatry, *48* (8), 803-812.
47. Manning-Courtney, P., Brown, J., Molloy, C. A., Reinhold, J., Murray, D., Sorensen-Burnworth, R., Messerschmidt, T., & Kent, B. (2003). Diagnosis and treatment of autism spectrum disorders. Current Problems in Pediatric and Adolescent Health Care, *33* (9), 283-304.
48. Masi, G., Cosenza, A., Mucci, M., & Broredani, P. (2003). A 3-year naturalistic study of 53 preschool children with PDD treated with Risperidone. Journal of Clinical Psychiatry, *64* (9), 1039-1046.
49. McCarthy, H. (2006). Survey of Children Adopted from Eastern Europe – The Need for Special School Services. Available online at: http://www.postadoptinfo.org/research/survey_results.php. Last viewed on October 25, 2013.
50. McConnell, S. (2002). Interventions to Facilitate Social Interaction for Young Children with Autism: Review of Available Research and Recommendations for Educational Intervention and Future Research. Journal of Autism and Developmental Disorders, *32* (5), 351-372.
51. McDougle, C. J., Erickson, C. A., Stigler, K. A., & Posey, D. J. (2005). Neurochemistry in the

BlueChoice HealthPlan-Medicaid UM Guideline

Subject: Autism Spectrum Disorder Services

Current Effective Date: 07/01/2017

Status: Active

Last Review Date: 06/11/2021

pathophysiology of autism. Journal of Clinical Psychiatry, 66 (Suppl. 10), 9-18.

52. Mostert, M. P. (2001). Facilitated communication since 1995; a review of published studies. Journal of Autism and Developmental Disorders, 31 (3), 287-313.
53. Myers, S. M. & Plauche' Johnson, C. (2007). Management of Children with Autism Spectrum Disorders. Pediatrics, 120 (5), 1162-1182.
54. Oneal, B. J., Reeb, R. N., Korte, J. R. & Butter, E. J. (2006). Assessment of Home-Based Behavior Modification Programs for Autistic Children: Reliability and Validity of the Behavioral Summarized Evaluation. Journal of Prevention & Intervention in the Community, 32 (1-2), 25-39.
55. Ospina, M. B., Krebs Seida, J., Clark B., Karkhaneh, M., Hartling, L., Tjosvold, L., Vandermeer, B. & Smith, V. (2008). Behavioural and Developmental Interventions for Autism Spectrum Disorder: A Clinical Systematic Review. PLoS ONE, 3 (11), e3775.
56. Ottenbacher, K. (1982). Sensory Integration Therapy: Affect or Effect. American Journal of Occupational Therapy, 36 (9), 571-578.
57. Posey, D. J. & McDougle, C. J. (2002). Autism: A three-step practical approach to making the diagnosis. Current Psychiatry, 1 (7), 20-28.
58. Reichow, B. & Wolery, M. (2008). Comprehensive Synthesis of Early Intensive Behavioral Interventions for Young Children with Autism Based on the UCLA Young Autism Project Model. Journal of Autism and Developmental Disorders, 39 (1), 23-41.
59. Remington, B., Hastings, R. P., Kovshoff, H., degli Espinosa, F., Jahr, E., Brown, T., ALSford, P., Lemaic, M. & Ward, N. (2007). Early Intensive Behavioral Intervention: Outcomes for Children With Autism and Their Parents After Two Years. American Journal on Mental Retardation, 112 (6), 418-438.
60. Rogers, S. and Vismara, L. (2008). Evidence-based Comprehensive Treatments for Early Autism. Journal of Clinical Child and Adolescent Psychology, 37 (1), 8-38.
61. Rogers, S. J. & Ozonoff, S. (2005). Annotation: What do we know about sensory dysfunction in autism? A critical review of the empirical evidence. Journal of Child Psychology and Psychiatry, 46 (12), 1255-1268.
62. Roley, S., Clark, G., Bissell, J. & Brayman, S. (2003). Applying Sensory Integration Framework in Educationally Related Occupational Therapy Practice (2003 Statement). American Journal of Occupational Therapy, 57 (6), 652-659.
63. Sacco, R., Militerni, R., Frolli, A., Bravaccio, C., Griti, A., Elia, M., Curatolo, P., Manzi, B., Trillo, S., Lenti, C., Saccani, M., Schneider, C., Melmed, R., Reichelt, K. L., Pascucci, T., Puglisi-Allegra, S. & Persico, A. M. (2007). Clinical, morphological, and biochemical correlates of head circumference in autism. Biological Psychiatry, 62 (9), 1038-1047.
64. Sallows, G. O. & Graupner, T. D. (2005). Intensive Behavioral Treatment for Children With Autism: Four-Year Outcome and Predictors. American Journal on Mental Retardation, 110 (6), 417-438.

BlueChoice HealthPlan-Medicaid UM Guideline

Subject: Autism Spectrum Disorder Services

Current Effective Date: 07/01/2017

Status: Active

Last Review Date: 06/11/2021

65. Sandler, A. (2005). Placebo Effects in Developmental Disabilities: Implications for Research and Practice. Mental Retardation and Developmental Disabilities, *11* (2), 164-170.
66. Scanhill, L. (2005). Diagnosis and evaluation of pervasive developmental disorders. Journal of Clinical Psychiatry, *66* (Suppl. 10), 19-25.
67. Schaaf, R. C. & Miller, L. J. (2005). Occupational Therapy Using a Sensory Integrative Approach for Children with Developmental Disabilities. Mental Retardation and Developmental Disabilities, *11* (2), 143-148.
68. Schaaf, R. C. & Nightlinger, K. M. (2007). Occupational therapy using a sensory integrative approach: A case study of effectiveness. American Journal of Occupational Therapy, *61* (2), 239-246.
69. Science 2.0. How Well Do Adopted Children Really Adjust To A New Language? Available at: http://www.science20.com/news_releases/how_well_do_adopted_children_really_adjust_new_language. Last viewed on October 25, 2013.
70. Shevell, M., Ashwal, S., Donley, D., Flint, J., Gingold, M., Hirtz, D., Majnemer, A., Noetzel, M. & Sheth, R. (2003). Practice parameter: Evaluation of the child with global developmental delay: Report of the Quality Standards Subcommittee of the American Academy of Neurology and The Practice Committee of the Child Neurology Society. Neurology, *60* (3), 367-380.
71. Shook, G. L., Ala'i-Rosales, S. & Glenn, S. (2002). Training and Certifying Behavior Analysts. Behavior Modification, *26* (1), 27-48.
72. Sikora, D. M., Hall, T. A., Hartley, S. L., Gerrard-Morris, A. E. & Cagle, S. (2008). Does Parent Report of Behavior Differ Across ADOS-G Classifications: Analysis of Scores from the CBCL and GARS. Journal of Autism and Developmental Disorders, *38* (3), 440-448.
73. Smith, S., A., Press, B., Koenig, K. P., & Kinnealey, M. (2005). Effects of sensory integration intervention on self-stimulating and self-injurious behaviors. American Journal of Occupational Therapy, *59*, 418-425.
74. Stahmer, A., Ingersoll, B., & Carter, C. (2003). Behavioral approaches to promoting play. Autism, *7* (4), 401-413.
75. South Carolina Department of Health and Human Services (SCDHHS). (2006). Application for 1915 HCBS Waiver. Available online at: <https://www.scdhhs.gov/sites/default/files/PDD%20Waiver%20Application.pdf>. Last viewed on October 25, 2013.
76. The University of Stavanger (2008, December 19). Up To A Third of Children Adopted To Norway From Abroad Are Having Problems With Language Proficiency. ScienceDaily. Available at: <http://www.sciencedaily.com/releases/2008/12/081219073053.htm>. Last viewed on October 25, 2013.
77. U.S. Department of Defense. (2007). Department of Defense Report and Plan on Services to

BlueChoice HealthPlan-Medicaid UM Guideline

Subject: Autism Spectrum Disorder Services

Current Effective Date: 07/01/2017

Status: Active

Last Review Date: 06/11/2021

Military Dependent Children with Autism. Available online at:

http://www.bacb.com/Downloadfiles/707_DoD_TRICARE_rpt.pdf. Last viewed on October 25, 2013.

78. Vargas, S. & Camilli, G. (1999). A meta-analysis of research on sensory integration treatment. American Journal of Occupational Therapy, *53* (2), 189-198.
79. Volkmar, F., Paul, R., Klin, A. & Cohen, D. (2005). Handbook of Autism and Pervasive Developmental Disorders (Vol. 1, 3rd ed., text rev.). Hoboken, NJ: John Wiley & Sons, Inc.
80. Warren, Z., Veenstra-VanderWeele, J., Stone, W., Bruzek, J. L., Nahmias, A. S., Foss-Feig, J. H., Jerome, R. N., Krishnaswami, S., Sathe, N. A., Glasser, A. M., Surawicz, T., McPheeters, M. L. (2011). Therapies for Children With Autism Spectrum Disorders. Comparative Effectiveness Review No. 26. (Prepared by the Vanderbilt Evidence-based Practice Center under Contract No. 290-2007-10065-I.) AHRQ Publication No. 11-EHC029-EF. Rockville, MD: Agency for Healthcare Research and Quality. April 2011. Available online at: http://www.effectivehealthcare.ahrq.gov/ehc/products/106/656/CER26_Autism_Report_04-14-2011.pdf. Last viewed on October 25, 2013.
81. Watling, R. L., Deitz, J. & White, O. (2001). Comparison of Sensory Profile scores of young children with and without autism spectrum disorders. American Journal of Occupational Therapy, *55* (4), 416-423.
82. Watling, R., Tomchek, S., & LaVesser, P. (2005). The scope of occupational therapy services for individuals with autism spectrum disorders across the lifespan. The American Journal of Occupational Therapy, *59* (6), 680-683.
83. Werner DeGrace, B. (2004). The everyday occupation of families with children with autism. American Journal of Occupational Therapy, *58*, 543-550.
84. Whalen, C. & Schreibman, L. (2003). Joint attention training for children with autism using behavior modification procedures. Journal of Child Psychology and Psychiatry, *44* (3), 456-468.
85. Windsor, J., Doyle, S. S., & Siegel, G. M. (1994). Language acquisition after mutism; a longitudinal case study of Autism. Journal of Speech and Hearing Research, *37* (1), 96-105.
86. Yoder, P. & McDuffie, A. (2006). Teaching young children with autism to talk. Seminars in Speech and Language, *27* (3), 161-172.
87. Yoder, P. & Stone, W. L. (2006). Randomized comparison of two communication interventions for preschoolers with autism spectrum disorders. Journal of Consulting and Clinical Psychology, *74* (3), 426-435.

Websites for Additional Information

South Carolina Department of Health and Human Services www.scdhhs.gov **Accessed August 9, 2023

BlueChoice HealthPlan-Medicaid UM Guideline

Subject: Autism Spectrum Disorder Services

Current Effective Date: 07/01/2017

Status: Active

Last Review Date: 06/11/2021

History

Status	Date	Action
New	5-3-2017	Created
Approved	6-5-2017	Medical Operations Committee, MOC
Updated	6/24/19	Updated ASD CPT codes
Reviewed and Approved	06/25/2020	Medical Operations Committee, MOC
Reviewed	06/11/2021	Reviewed-no changes

BlueChoice HealthPlan-Medicaid UM Guideline

Subject: Autism Spectrum Disorder Services
Status: Active

Current Effective Date: 07/01/2017
Last Review Date: 06/11/2021

Approved	06/17/2021	Medical Operations Committee (MOC)

This page includes links to third party websites. Those organizations are solely responsible for the contents and privacy policies on its site.